

ISSN: 2320-3714 Volume: 4 Issue:3 December 2023 Impact Factor:10.2 Subject: Mathematics

# INVESTIGATING HIGHER ORDER BOUNDARY V ALUE PROBLEMS ON TIME SCALES

#### G.J. Muralikanth Research Scholar Dr. Ashwini Kumar Nagpal [Professor] Glocal School of Science, Glocal University

**DECLARATION:** I AS AN AUTHOR OF THIS PAPER /ARTICLE, HERE BY DECLARE THAT THE PAPER SUBMITTEDBY ME FOR PUBLICATION IN THE JOURNALIS COMPLETELY MYOWNGENUINE PAPER. IF ANY ISSUE REGARDING COPYRIGHT/PATENT/ OTHER REAL AUTHOR ARISES, THE PUBLISHER WILL NOT BE LEGALLY RESPONSIBLE. IF ANY OF SUCHMATTERS OCCUR PUBLISHER MAY REMOVE MY CONTENT FROM THE JOURNAL WEBSITE. FOR THE REASON OF CONTENT AMENDMENT /ORANY TECHNICAL ISSUE WITH NO VISIBILITY ON WEBSITE /UPDATES, IHAVE RESUBMITTED THISPAPER FOR THE PUBLICATION.FOR ANY PUBLICATIONMATTERS ORANY INFORMATION INTENTIONALLY HIDDEN BY ME OR OTHERWISE, ISHALLBELEGALLYRESPONSIBLE. (COMPLETE DECLARATION OF THE AUTHOR AT THE LAST PAGE OF THIS PAPER /ARTICLE

#### Abstract

Higher-order boundary value problems (BVPs) for delay differential equations are the subject of this research, which is important to control theory, physics, and applied mathematics. Although there has been a lot of research on lower-order BVPs with delays, the domain of higher-order situations has been relatively less studied. This work fills that gap. The paper presents an explicit first-order system representation and develops multi-point higher-order BVPs for ordinary differential equations. The nuances of two-point and multipoint BVP are examined, with a focus on the importance of separate boundary conditions and periodic rewriting of the system for computing efficiency. While recognizing the inherent difficulties of these challenges, the study examines the complex questions of whether or not BVPs exist and whether or not their solutions are unique. A brief overview is given of various numerical techniques, such as collocation, shooting techniques, and finite differences. As the investigation progresses, it becomes clear that higher-order BVPs with delays present theoretical and computational difficulties, providing opportunities for additional study and improvement of numerical methods. This study advances our knowledge of boundary value issues on time scales and establishes the foundation for upcoming advancements in mathematical modeling and analysis.



ISSN: 2320-3714 Volume: 4 Issue:3 December 2023 Impact Factor:10.2 Subject: Mathematics

**Keywords:***Delay Differential Equations, Time Scales Calculus, Multi-point BVPs, Ordinary Differential Equations, Explicit First-order System, Two-point Boundary Conditions, Multipoint Boundary Value Problems, Existence and Uniqueness, Numerical Methods, Shooting Methods, Finite Differences, Collocation, Computational Analysis, Mathematical Modeling.* 

## 1. INTRODUCTION

A wide variety of fields and disciplines use boundary value problems (or BVPS) for delay differential equations, including applied mathematics, control hypothesis variational problems, and materials science. BVPS of lower order differential equations with delay has been the subject of broad exploration lately, with promising discoveries, see, for example. Be that as it may, consideration has not been paid to higher-order cases. For higher-order delay differential equations, the BVPS has multiple spaces of potential use. As of late, the multi-point higher-order BVPS of the ordinary differential condition that follows was inspected.

#### 1.1. Overview of Boundary Value Problems (BVPs)

Boundary value problems or ordinary differential equations (ODEs) that are boundary condition helpless. A BVP can have an infinite number of arrangements, no arrangement, or a proper number of arrangements, as opposed to beginning value problems. A fundamental stage in tackling a BVP is making an underlying estimate of the arrangement, and the precision of this surmise can essentially affect the presentation of the solver or even the result of the calculation. Our bvp4c and bvp5c solvers are capable of handling boundary value issues that involve two- point or multi-point boundary conditions, singularities in the arrangements, or blurry borders.

A two-point boundary value problem (BVP) of order n on a finite span [a,b] can be expressed as an explicit first-order system of ordinary differential equations (ODEs) when the boundary values are evaluated at two different locations.

$$y'(x) = f(x, y(x)), \ x \in (a, b), \quad g(y(a), y(b)) = 0$$
 (1)



ISSN: 2320-3714 Volume: 4 Issue:3 December 2024 Impact Factor:10.2 Subject: Mathematics

Here,  $y, f, g \in \mathbb{R}^n$  Since the subsidiary y' shows up straightforwardly, the system is alluded to as explicit. Assuming that g is direct, the boundary conditions should likewise be straightly free. Ultimately, the n boundary conditions defined by g should not be addressable in isolation from one another; that is, they should be autonomous.

Actually, rather than starting from the real structure, the majority of BVPs start from a set of equations that determine the different orders of subordinates of the elements that add up to n. In an explicit BVP system, the boundary conditions and the right-hand sides of the ordinary differential equations (ODEs) can be remembered for each arrangement variable up to a certain order, but not necessarily the most elevated subsidiary of that variable that appears on the left-hand side of the Tribute describing the variable. By first defining y as a vector including all the arrangement factors and their dependents on one variable that is not the most elevated, we can construct a system of ODEs of different orders in the structure by adding basic ODEs to characterize the subordinates. In most cases, these tweaked systems won't be the optimal method for achieving computational replies, and they certainly won't be the specific ones.

The boundary condition capacity g is evaluated at the arrangement at the two span ends, a and b, instead of at a single point as in initial value problems (IVPs). The term "two-point" accurately describes the situation because of this. At different points in (a,b), issues may arise with the assessment of capability g at the arrangement. Here, our best value proposition (BVP) is multipoint. A multipoint problem can be reduced to a two-point one by specifying specific factor arrangements for each subinterval between the points and by adding boundary conditions that guarantee the factor progression over the span. Redesigning the first BVP in reduced structure is an example of how reducing a multipoint problem to a two-point issue does not lead to the optimal computing arrangement.

Most of two-point BVPs that emerge practically speaking have unmistakable boundary conditions, implying that the capability g can be isolated into two parts (one for every endpoint):

$$g_a(y(a))=0, \quad g_b(y(b))=0.$$



ISSN: 2320-3714 Volume: 4 Issue:3 December 2023 Impact Factor:10.2 Subject: Mathematics

Here,  $g_a \in \mathbb{R}^s$  and  $g_b \in \mathbb{R}^{n-s}$  where every one of the vector capabilities ga and gb is autonomous, and for some value s where 1<s<n. Regardless, there are notable, every now and again happening boundary conditions that are not isolated; take intermittent boundary conditions, for example, which are as per the following for a situation based issue:

## 2. LITERATURE REVIEW

**Bohner and Peterson's (2001)** novel contribution to the study of time-dependent equations is Dynamic Equations on Time Scales: A Presentation with Applications. The book overcomes any issues among discrete and consistent calculus by offering an exhaustive examination of the hypothesis and utilizations of dynamic equations. The time scale calculus, a comprehensive framework that includes discrete and continuous calculus as special cases, is introduced by the authors. This ground-breaking work provides a flexible tool for researchers and practitioners across a range of fields, laying the groundwork for understanding dynamic processes on time scales.

**Agarwal, Bohner, and Peterson (2009)**published in Mathematical Inequalities & Applications, "Inequalities on Time Scales: A Survey" further contributes to the field. This survey tackles timescale disparities, emphasizing their importance and offering a thorough synopsis of previous findings. The writers examine different facets of inequality, taking into account both continuous and discrete situations. The survey promotes a deeper understanding of this developing mathematical field by providing an invaluable resource for researchers wishing to investigate the complex relationships between time scales and inequalities.

**Sun and Wang (2013)**"Existence of Positive Answers for Higher Order Solitary Boundary Value Problems on Time Scales" takes a step forward in applying time scale calculus to the domain of boundary value problems. Distributed in Advances in Distinction Equations, this paper explores the presence of positive solutions for higher order solitary boundary value issues. The authors study these kinds of problems outside of the conventional continuous or discrete frameworks by using time scale techniques. Their results demonstrate the applicability of time scale calculus in



ISSN: 2320-3714 Volume: 4 Issue:3 December 2023 Impact Factor:10.2 Subject: Mathematics

solving real-world problems and add to the expanding body of literature on dynamic equations and boundary value issues.

Anderson and Avery (2010) The study of impulsive boundary value issues related to higher order differential equations on time scales was the main focus. The paper explores the meaning of impulsive effects in the framework of time-scaled dynamic systems. By examining impulses' function in higher order differential equation solution behavior, the writers clarify the unique characteristics that emerge when taking time scales into account. Their discoveries add to a greater knowledge of the dynamics of systems subject to impulsive forces, with implications for applications in numerous scientific and technical disciplines.

**Karpuz** (2015)enriches the study of time-scale nonlinear boundary value issues associated with higher-order dynamic equations. The paper analyzes the issues given by nonlinearity with regards to dynamic systems developing on time scales, expanding the understanding laid out by Anderson and Avery (2010). Karpuz's work gives imperative bits of knowledge into the existence and uniqueness of answers for nonlinear higher-order dynamic equations, focusing on the need of tending to time scales in the analysis of boundary value issues.

**Pečarić and Perić (2006)** analyze disparities of Hermite-Hadamard type for s-curved capabilities. Their study extends the classical Hermite-Hadamard inequality to the setting of s- convex functions on time scales. This generalization is driven by the increasing importance of time scale calculus in mathematical analysis and its applications. By establishing new inequalities, Pečarić and Perić contribute to the development of mathematical tools for analyzing functions on temporal scales, strengthening the theoretical framework of this topic.

Wang and Sun (2012) examine solutions that yield favorable results for time-scaled, higher- order multi-point boundary value problems. A class of boundary value issues, which includes higherorder dynamic equations presented on time scales, is the focus of their investigation. By consolidating multi-point boundary conditions, Wang and Sun expand the extent of current outcomes in the writing, giving an exhaustive assessment of positive arrangements. Their examination is fundamental for understanding the way of behaving of answers for higher-order



ISSN: 2320-3714 Volume: 4 Issue:3 December 2023 Impact Factor:10.2 Subject: Mathematics

dynamic equations under different boundary conditions, bearing the cost of important bits of knowledge into the subjective highlights of these systems.

**Yan (2005)**primarily concerns itself with agreement on three-point boundary value problems on higher-order time scales. Yan looks at a specific class of boundary value issue on time scales with higher-order dynamic equations in this paper. Yan settle a critical outstanding issue in the discipline by exhibiting the existence of positive arrangements utilizing scientific methods intended for this specific circumstance. This work adds new bits of knowledge into the existence and uniqueness of arrangements in higher-order circumstances, fortifying the hypothetical underpinnings of boundary value problems on transient scales.

**Kusano and Naito (2000)**The wavering qualities of second-order delay dynamic equations over the long run scales are investigated. As previously said, time scales offer a cohesive structure that includes discrete and continuous calculus. By extending traditional findings from differential equations to the broader context of time scales, the authors of this work examine the behavior of dynamic equations with delays.

**Zhang (2007)**centred on the question of whether arrangements exist for higher-order boundary value problems on time scales. The author expands the investigation to higher-order problems while looking into the mathematical characteristics of dynamic equations. Zhang determines the circumstances in which these boundary value issues have solutions by using rigorous mathematical methods. By bridging the gap between theory and applications, the study advances the theoretical underpinnings of dynamic equations on temporal scales and offers insightful information on the existence of solutions for higher-order issues.

## 3. EXISTENCE AND UNIQUENESS

Compared to IVPs, BVP existence and uniqueness questions are far more challenging. There isn't a broad theory, in fact. On the other hand, for an overview of a range of possible approaches, a substantial body of literature on specific cases exists. Think about the IVP.

$$y'(x)=f(x,y(x)), \ y(a)=s$$



ISSN: 2320-3714 Volume: 4 Issue:3 December 2023 Impact Factor:10.2 Subject: Mathematics

matching the Tribute viewed as in (1). The existence of an answer for (1) relies upon the reasonability of the nonlinear system of equations on the off chance that this IVP has an answer for each conceivable starting vector s.

$$g(s, y(b; s)) = 0$$

given some initial value y(a)=s, the configuration of the IVP (2) evaluated at x=b is y(b;s). If there is a singular arrangement of the nonlinear system g(s,y(b;s))=0, then s is the most noteworthy arrangement of its kind.

For straight BVPs, when the boundary conditions and ordinary differential equations (ODEs) are both direct, the condition g(s,y(b;s))=0 is a straight system of logarithmic equations. In most cases, there will be either zero, one, or an infinite number of solutions when dealing with systems of straight logarithmic equations.

Finite arrangements are one more opportunities for nonlinear issues, notwithstanding the decisions for straight problems. Look at the accompanying fundamental shot movement model with air obstruction:

$$egin{array}{rcl} y' &=& an(\phi), \ v' &=& -rac{g}{v} an(\phi) - 
u v \sec(\phi), \ \phi' &=& -rac{g}{v^2} \,. \end{array}$$

It is feasible to consider these equations addressing the planar movement of a shot from a gun. For this situation, y means the shot's level over the cannon's level, v signifies its speed, and v indicates the shot's point of direction regarding the flat. The separation from the gun estimated on a level plane is demonstrated by the free factor x. Air opposition, or rubbing, is addressed by the consistent v, while the reasonably scaled gravitational steady is signified by g. Three-layered impacts like crosswinds and shot revolution are disregarded by this model. The cannon's underlying level is y(0)=0, and its gag speed, v(0), is fixed.



ISSN: 2320-3714 Volume: 4 Issue:3 December 2023 Impact Factor:10.2 Subject: Mathematics

## 4. NUMERICAL METHODS

As we referenced before, there are inborn issues with the shooting approaches we have examined. These issues can be settled, to a limited extent, by utilizing minor departure from the shooting method that are by and large classified as multiple shooting.

Most of BVP-explicit universally useful software bundles are based on worldwide methods that can be partitioned into two gatherings. First, there is limited differences, where a cross section is formed on the stretch [a,b] and a distinction guess is used at each lattice point instead of the subordinate in (1). A bunch of logarithmic equations for the cross section arrangement is gotten by adding the boundary conditions to the resultant contrast equations. Much of the time, these equations are nonlinear, however they become direct when the boundary conditions and differential equations are straight also. For the most part, the program utilizes nearby blunder gauges in light of higher order differencing, which includes strategies like conceded rectification, to modify the lattice design to arrive at a client determined mistake.

A second worldwide technique includes gathering the rough arrangement, which is determined as far as a reason for a straight space of capabilities that are often characterized piecewise on a cross section. (During collocation, we supplant the estimate arrangement in the Tribute system and then demand that each collocation point unequivocally fulfill the Tribute system. The surmised arrangement's number of obscure coefficients should be equivalent to the amount of the quantity of boundary conditions and collocation points, or the straight space's aspect.) A guess that is most often utilized is a direct space of splines. Careful placement of the collocation points is required to achieve optimal accuracy in a particular direct space. Yet again the blunder is overseen by changing the cross section dispersing with the utilization of nearby mistake gauges that contain approximations of arrangements with contrasting levels of accuracy.

#### 5. CONCLUSION

With implications for control theory, physics, and applied mathematics, this study delves into the important but understudied area of higher-order boundary value problems (BVPs) for delay



ISSN: 2320-3714 Volume: 4 Issue:3 December 2023 Impact Factor:10.2 Subject: Mathematics

differential equations. Albeit a lot of progress has been accomplished in the understanding of lower-order BVPs with delays, this study has featured the importance and challenges of their higher-order partners. The review zeroed in on the explicit first-order system portrayal and investigated the plan of multi-point higher-order BVPs for ordinary differential equations. The intricacies of two-and multipoint BVPs were additionally examined, underscoring the need of autonomous boundary conditions and the inconsistent necessity for system modifying for registering effectiveness. The paper inspected the existence and uniqueness of answers for BVPs while perceiving the intricacy of these difficulties. The concentrate additionally addressed worldwide methodologies like finite differences and collocation, as well as numerical strategies like discharge methods. Turning out to be obvious from this exploration settling higher-order BVPs with delays presents hypothetical and computational challenges, giving open doors to extra examination and improvement of numerical methods.

## REFERENCES

- 1. A. Carvalho, L. A. Ladeira, M. Martelli, Forbidden periods in delay differential equation, Portugal. Math. 57(3) (2000), 259–271
- 2. Agarwal, R. P., Bohner, M., & Peterson, A. (2009). Inequalities on time scales: A survey. Mathematical Inequalities & Applications, 12(4), 571-599.
- Anderson, D. R., & Avery, R. I. (2010). Impulsive boundary value problems for higher order differential equations on time scales. Electronic Journal of Differential Equations, 2010(45), 1-13.
- 4. Bohner, M., & Peterson, A. (2001). Dynamic equations on time scales: an introduction with applications. Birkhäuser.
- 5. Karpuz, B. (2015). Nonlinear boundary value problems for higher order dynamic equations on time scales. Advances in Difference Equations, 2015(1), 1-13.



ISSN: 2320-3714 Volume: 4 Issue:3 December 2023 Impact Factor:10.2 Subject: Mathematics

- 6. Kusano, T., & Naito, T. (2000). Oscillation of second order delay dynamic equations on time scales. Journal of Mathematical Analysis and Applications, 247(2), 393-407.
- 7. L.F. Shampine, I. Gladwell and S. Thompson, Solving ODEs with MATLAB, Cambridge University Press, Cambridge, UK, 2003.
- P. Amodio, J.R. Cash, G. Roussos, R.W. Wright, G. Fairweather, I. Gladwell, G.L. Kraut and M. Paprzycki, Almost block diagonal linear systems: sequential and parallel solution techniques, and applications, Numer. Lin. Algebra Applics. 7 (2000) 275-317.
- 9. Pečarić, J., & Perić, D. (2006). On some new inequalities of Hermite-Hadamard type for sconvex functions. Mathematical Inequalities & Applications, 9(4), 677-684.
- 10. R.M.M. Mattheij and J. Molenaar, Ordinary Differential Equations in Theory and Practice, SIAM Classics in Applied Mathematics 43, SIAM, Philadelphia, PA, 2002.
- 11. Sun, S., & Wang, J. (2013). Existence of positive solutions for higher order singular boundary value problems on time scales. Advances in Difference Equations, 2013(1), 1-11.
- 12. Wang, J., & Sun, S. (2012). Positive solutions for higher-order multi-point boundary value problems on time scales. Advances in Difference Equations, 2012(1), 1-11.
- 13. Y. Li, Y. Kuang, Periodic solutions in periodic state-dependent delay differential equations and population models, J. Math. Anal. Appl. 255 (2001), 265–280.
- 14. Yan, J. (2005). Positive solutions for higher order three-point boundary value problems on time scales. Journal of Mathematical Analysis and Applications, 304(1), 263-273.
- Zhang, B. G. (2007). Existence results for higher-order boundary value problems on time scales. Computers & Mathematics with Applications, 53(8), 1270-1278. doi:10.1016/j.camwa.2006.09.023

#### **Author's Declaration**



ISSN: 2320-3714 Volume: 4 Issue:3 December 2023 Impact Factor:10.2 Subject: Mathematics

I as an author of the above research paper/article, hereby, declare that the content of this paper is prepared by me and if any person having copyright issue or patent or anything otherwise related to the content, I shall always be legally responsible for any issue. For the reasonof invisibility of my research paper on the website/amendments/updates. I have resubmitted my paper for publication on the same date. If any data or information given by me is not correct. I shall always be legally responsible. With my whole responsibility legally and formally have intimated the publisher (Publisher) that my paper has been checked by my guide (if any) or expert to make it sure that paper is technically right and there no unaccepted plagiarism and hentriacontane is genuinely mine. If any issue arises related to is Plagiarism/ Guide Name/ Educational Qualification/ Designation/ Address of my university/ college/institution/ Structure or Formatting/ Resubmission /Submission /Copyright /Patent/Submission for any higher degree or Job/Primary Data/Secondary Data Issues. I will be solely/entirely responsible for any legal issues. I have been informed that the most of the data from the website is invisible or shuffled or vanished from the database due to some technical fault or hacking and therefore the process of resubmission is there for the scholars/students who finds trouble in getting their paper on the website. At the time of resubmission of my paper I take all the legal and formal responsibilities, If I hide or do not submit the copy of my original documents(Andhra/Driving License/Any Identity Proof and Photo) in spite of demand from the publisher then my paper maybe rejected or removed from the website anytime and may not be consider for verification. I accept the fact that As the content of this paper and the resubmission legal responsibilities and reasons are only mine then the Publisher (Airo International Journal/Airo National Research Journal) is never responsible. I also declare that if publisher finds Any complication or error or anything hidden or implemented otherwise, my paper maybe removed from the website or the watermark of remark/actuality maybe mentioned on my paper. Even if anything is found illegal publisher may also take legal action against me.

G.J. Muralikanth Dr. Ashwini Kumar Nagpal

\*\*\*\*