

OXIDATIVE BEHAVIOR OF PHENYLALANINE IN AN ACIDIC DMF–WATER MEDIUM A KINETIC STUDY WITH PYRIDINIUM CHLOROCHROMAT

Shobhag Singh Sisodiya

Research Scholar

Chemistry

Dr. Rama Nand Yadav

(Professor)

Department of Science, Sunrise University Alwar

DECLARATION: I AS AN AUTHOR OF THIS PAPER /ARTICLE, HERE BY DECLARE THAT THE PAPER SUBMITTED BY ME FOR PUBLICATION IN THE JOURNAL IS COMPLETELY MY OWN GENUINE PAPER. IF ANY ISSUE REGARDING COPYRIGHT/PATENT/ OTHER REAL AUTHOR ARISES, THE PUBLISHER WILL NOT BE LEGALLY RESPONSIBLE. IF ANY OF SUCH MATTERS OCCUR PUBLISHER MAY REMOVE MY CONTENT FROM THE JOURNAL WEBSITE. FOR THE REASON OF CONTENT AMENDMENT /OR ANY TECHNICAL ISSUE WITH NO VISIBILITY ON WEBSITE /UPDATES, I HAVE RESUBMITTED THIS PAPER FOR THE PUBLICATION. FOR ANY PUBLICATION MATTERS OR ANY INFORMATION INTENTIONALLY HIDDEN BY ME OR OTHERWISE, I SHALL BE LEGALLY RESPONSIBLE. (COMPLETE DECLARATION OF THE AUTHOR AT THE LAST PAGE OF THIS PAPER /ARTICLE)

Abstract

In a watery DMF model including perchloric acid, studies on the oxidation of phenyl alanine by pyridinium chlorochromate have been completed. It was observed that the initial sales contact with [PCC] had the quickest response time. In a DMF-water (70:30%) mixture under perchloric acid, the oxidation of phenyl alanine by pyridinium chlorochromate generates the energy necessary to produce the corresponding aldehyde. First requesting responses happen in [PCC], [HClO₄], and [AA]. Phenyl alanine showed energy of the Michaelis-Menten type. At different temperature, the response rates were found, and as far as possible were joined up. Acrylonitrile doesn't go through polymerization considering the response. The rate increases as the social event of DMF in the fluid blend increments. A fitting system was proposed to understand the response.

Keywords: Oxidation Behaviour, Phenyl Alanine, Acidic Dimethyl Formamide (DMF), Kinetic, Water Medium, Pyridinium Chlorochromate (PCC).

1. INTRODUCTION

Our review's principal objective is to expand how we might interpret how a few amino acids oxidize when presented to PCC. The article portrayed the kinetics of pyridinium

chlorochromate's oxidation of methionine, oximes, unsaturated acids, cysteine, and alcohols. The oxidative contrast in glycine and alanine, l-isoleucine and nor-leucine, and phenylalanine by PDC in an aquo-acidic disastrous media was contemplated capably and recklessly. The oxidation of alanine by PCC in DMF-Water medium with perchloric horrendous present was addressed. Clearly no reports exist about PCC's oxidation of L-isoleucine and L-valine. Our review of the composing shows that not much assessment has been done on the oxidation of amino acids by various oxidants; we are especially curious to get to know the arrangement of PCC's oxidation of amino acids in acidic DMF-Water conditions.

Pyridinium chlorochromate is an extraordinarily versatile oxidizing expert that is immediately open. The making contains various evaluations on the oxidation of different substrates by pyridinium chlorochromate (PCC), including alcohols, methionine, oximes, unsaturated acids, and cysteine. Elucidates the oxidation of tyrosine by pyridinium chlorochromate don't seem to exist. Organosulfur compounds, for example, phenyl alanine (otherwise called thiocarbamide or sulfuric), have the equation $\text{H}_2\text{NSCNHPh}$. While it imparts underlying likenesses to urea (except for a sulphur iota instead of the oxygen particle), phenyl alanine is significantly unique in relation to urea regarding its qualities. One reagent utilized in natural union is phenyl alanine. A huge class of mixtures having the general construction $(\text{R}_1\text{R}_2\text{N})(\text{R}_3\text{R}_4\text{N})\text{C}=\text{S}$ are alluded to be "phenyl alanine's." Thioamides and phenyl alanine's are connected.

1.1. Overview of Phenylalanine

An essential amino acid called phenylalanine is needed to synthesise L-aspartame, an artificial sweetener used by diabetics that is 200 times sweeter than sugar. It is produced in 15,000 tonnes annually at an approximate cost of 35 euros per kilogramme on the global market, and it is utilised in a wide variety of meals and drinks. Aspartame is currently manufactured via a biological procedure, as opposed to its original chemical manufacturing method. L-phenylalanine and L-aspartic acid, the two most crucial components of L-aspartame, are generated via fermentation and biocatalysts, respectively. The Holland Sweetener Company specifically links these two amino acids together using the bacterial enzyme thermolyzing.

1.2. Pyridinium Chlorochromate

The formula for pyridinium chlorochromate (PCC) is $[C_5H_5NH]^+[CrO_3Cl]^-$. It is a yellow-orange salt. This reagent is mostly employed in organic synthesis to generate carbonyls by oxidising alcohols. Many related chemicals with comparable reactivity are known. Compared to many other reagents, PCC has the advantage of selectively oxidising alcohols to aldehydes or ketones.

One bar up the oxidation stepping stool, from essential alcohols to aldehydes and from optional alcohols to ketones, is where PCC oxidizes alcohols. PCC doesn't oxidize aldehydes to carboxylic acids, as opposed to chromic corrosive. Similar to or indistinguishable from: Essential alcohols can likewise be oxidized to aldehydes by utilizing CrO_3 and pyridine, the Collins reagent. The following are two occasions where PCC is being used.

- The oxidised version of either of these alcohols will be created if one equivalent of PCC is added. The by-products are pyridinium hydrochloride and Cr (IV), which are shown in grey.
- How much water in the response should be painstakingly thought of. On the off chance that water is accessible, it can respond with the aldehyde to shape the hydrate, which can then be additionally oxidized by the presence of one more likeness PCC. With ketones, this isn't an issue since H isn't associated straightforwardly to C.

1.3. Dimethyl Formamide (DMF)

DMF is primarily used as a low-evaporation-rate solvent. Acrylamide is utilised in the manufacturing of polymers and acrylic fibres. In the creation and assembling of bug sprays, cements, engineered cowhides, filaments, movies, and surface coatings, in addition to other things, it fills in as a dissolvable in peptide coupling for drugs.

- It is utilised as a reagent in the Vilsmeier-Haack reaction, another helpful process for generating aldehydes, as well as in the Bouveault aldehyde synthesis.
- The Heck reaction frequently uses it as a solvent.
- It is a frequently used catalyst in the production of acyl halides, specifically oxalyl or thionyl chloride-based acyl chlorides from carboxylic acids. The reversible production of

imidoyl chloride, popularly referred to as the "Vilsmeier reagent," is the catalytic mechanism.

2. RESEARCH METHODOLOGY

2.1. Materials

In the momentum review, twofold refined water was used to set up the arrangements and phenyl alanine scientific grade (SRL) was utilized as provided (not set in stone by its dissolving point). A known volume of corrosive was weakened in water to make perchloric corrosive. Involving phenolphthalein as a marker, a titration with recently normalized sodium hydroxide was utilized to normalize the created arrangement. The dissolvable source was insightful grade DMF (SRL) that had gone through twofold refining. Also, any remaining synthetic substances used were 99.9% unadulterated insightful grade reagents.

2.2. Pyridinium Chlorochromate Preparation

Utilizing a framework illustrated in the creation, pyridinium chlorochromate was made in the investigation office. 100 g of CrO_3 was immediately blended into 184 ml of 6 M hydrochloric horrendous. The strong might be set something to the side for wide time frames at room temperature without changing considering how it isn't perceivably hygroscopic. Iodometric testing and UV-Vis and IR range evaluation were utilized to avow the perfection.




Figure 1:Planning of Pyridinium Chlorochromate


2.3. Method

At 313 K, the response was led in a DMF-water [70% (v/v) DMF] dissolvable arrangement under pseudo first request conditions. The reaction was started by merging phenyl alanine and PCC in a thermally equilibrated plan that moreover contained the essential proportions of perchloric destructive. According to the JASCO model 7800 UV/Vis spectrophotometer, the absorbance of PCC at 354 nm in a 1 cm cell placed in the thermostatic compartment reportedly showed how much had diminished after the response.

Over 70% of the response was finished during the motor runs, and great first request kinetics were seen. The incline of the log(absorbance) against time plot was utilized to ascertain pseudo- first request rate constants, or k_{obs} .

2.4. Analysis of products and stoichiometry

In a 70% vol. DMF, 30% vol. water mix (v/v), the following conditions were used for the assessment: $[PCC] > [phenyl\ alanine]$, 0.3 mol/dm³ [HClO₄], and it was left to remain at 40 C. The evaluation was driven at 313 K. This drew in for the affirmation of the response's stoichiometry. Following the completion of the response, the absorbance at 354 nm was utilized to quantify the PCC focus. Under powerful conditions, the abstract thing study was coordinated. The 2,4-dinitrophenylhydrazine auxiliary of aldehyde thought about the separating verification of the fundamental reaction thing. The ammonium molecule and carbon dioxide were perceived using Nessler's reagent test and the lime water test, independently. After the reaction was finished, Cr (III) was checked by the reaction game plan's UV-perceptible spectra. One possible depiction of the saw stoichiometry is:



3. DATA ANALYSIS

3.1. The stability of chlorochromate produced from pyridinium

The strategy of PCC in DMF-water, which contains 70% (v/v) DMF, is to combine Lambert's Rule at 354 nm. The optical thickness and spectra of the PCC plan were unchanged upon wide standing or warming up to 335-345 K, particularly in the DMF-water mix containing 70% (v/v) DMF, when the substrate was not present.

3.2. This is the impact of PCC

For a constant concentration of $[\text{HClO}_4]$, temperature, and [amino acid] (phenyl alanine) [PCC] (the principal solicitation), the rate was directly proportional to the logarithm of [PCC] versus time. After adjusting the basic union of PCC, no discernible difference in the rate steady kobs could be seen.

3.3. The influence of the substrate

While the affiliation of PCC, the social occasion of hydrogen particles, and the temperature remained consistent, the reaction rate increased when the concentration of phenyl alanine increased from $1.2 \cdot 10^3$ to $5.0 \cdot 10^1$ mol/dm² (Table 1). This demonstrates that [phenyl alanine] is crucial to the rate in the core sales. Anyway, no matter what the way that the catch respect $[K_m = 0.004]$ is incredibly low, the worth shows the improvement of a perplexed that might be extraordinarily responsive, and that recommends that the middle will be uncommonly low at some unpredictable time. It has been discovered that the oxidation of a-amino damaging by Cr (VI) has a similar signature, which has been accepted.

Table 1: Change in pace in response to concentrations of perchloric acid, phenyl alanine, and PCC

$10^3 \times [\text{PCC}] \text{ mol/dm}^3$	$10^2 \times [\text{Amino acid}] \text{ mol/dm}^3$	$10 \times [\text{H}^+] \text{ mol/dm}^3$	$k_{\text{obs}} \times 10^5 \text{ s}^{-1}$
2.4	2.1	2.9	24.92
2.24	2.1	2.9	25.25

2.1	2.1	2.9	25.30
1.76	2.1	2.9	25.90
1.4	2.1	2.9	25.90
1.1	2.1	2.9	25.30
2.1	1.1	2.9	15.69
2.1	1.5	2.9	21.85
2.1	2.1	2.9	25.70
2.1	2.5	2.9	29.77
2.1	2.9	2.9	34.55
2.1	3.65	2.9	41.05
2.1	2.1	1.0	6.50
2.1	2.1	2.6	19.55
2.1	2.1	3.1	25.70
2.1	2.1	3.6	31.45
2.1	2.1	5.1	47.20
2.1	2.1	7.1	80.60

T = 313 K DMF = 70% (v/v).




Figure 2:Change in pace in response to concentrations of perchloric acid, phenyl alanine, and PCC

Table 2:Phenylalanine rate variation at 313 K with sodium sulphate fixing

$[\text{Na}_2\text{SO}_4] \times 10^3$ mol/dm^3	1.1	2.9	4.9	7.1	11.1
$10^5 k_{\text{obs}} (\text{s}^{-1})$	25.25	24.94	25.40	26.40	25.40




Figure 3:Phenylalanine rate variation at 313 K with sodium sulphate fixing

3.4. Ionic Strength's Impact

By changing the assembly of sodium sulphate, the impact of ionic strength was investigated. With the associations of phenyl alanine, PCC, and HClO_4 remaining unsurprising, the ionic strength in the response medium was changed some spot in the extent of 1.0 and $11.0 \cdot 10^3 \text{ mol/dm}^3$. Any remaining boundaries were kept steady. It has been noticed that the ionic strength perceptibly affects the rate. This recommends that an iron and a nonpartisan particle or two unbiased atoms might be engaged with the response.

3.5. The composition of the solvent

At the point when the DMF (% v/v) in the response blend was changed while keeping up with different boundaries steady, it was found that the response rate was essentially affected. As the volume level of DMF expanded, the pace of response additionally expanded. The effect of the dielectric constant (D) of the medium on the rate of reactions in the liquid state has been the subject of several quantitative explanation proposals. In a quick plot of $\log k_{\text{obs}}$ versus $1/D$, Amis demonstrated that a positive trend indicates a positive particle dipole response, whereas a negative trend indicates the responsibility of two dipoles or a negative atom dipole response, limiting the illustration to a particle dipole structure or a non-point of approach between the two.

Table 3: Changes in rate corresponding to dissolvable creation

DMF: H ₂ O	$k_{\text{obs}} \times 10^5 \text{ s}^{-1}$
70:30	25.90
60:40	14.41
50:50	8.003
40:60	6.17
30:70	5.012
20:80	2.23




Figure 4:Changes in rate corresponding to dissolvable creation

Table 4:Change in rate with temperature

Temperature (k)	299	302	307	312	317	322
$k_{obs} \times 10^5 \text{ s}^{-1}$	8.75	12.65	16.84	25.70	30.14	43.80




Figure 5: Change in rate with temperature

3.6. The impact of temperature

It was found that when temperature expanded, the response's rate consistent expanded (Table 4). The plot of $\log k$ versus $1/T$ yielded the activation energy, from which the initiation boundaries were determined (Table 5). As would be anticipated for a bimolecular interaction, the entropy of enactment is negative. During the rate-picking step, a non-cyclic reactant can be improved into a cyclic impermanent by using the negative worth additionally.

Table 5:Boundaries of actuation

Parameters	E_a (kJ mol ⁻¹)	ΔH^\ddagger (kJ mol ⁻¹)	ΔS^\ddagger (kJ mol ⁻¹)	ΔF^\ddagger (Jk1 mol ⁻¹)
Values	53.60	51.00	-93.763	80.33

Plotting the converse rate steady against the backwards substrate fixation (phenyl alanine) gives proof to the mind-boggling arrangement. It has been noted that a tiny and negative entropy of activation will cause a slowdown in the reaction.

3.7. Acrylonitrile and pyridine's effects

Since there is neither a rate diminishes within the sight of without stabilizer acrylonitrile nor a smooth appearance under motor circumstances, the inclusion of extremist cycle is precluded. At the point when pyridine is added, the pace of response stays unaltered, proposing that PCC is steady and not hydrolysed under the explored conditions.

3.8. Perchloric acid's effects

It was found that the rate expanded as the amount of perchloric corrosive expanded, considerably under fixed convergences of PCC and phenyl alanine and under other steady circumstances (Table 1).

Since the anion structure focus will be exceptionally low under the ongoing trial conditions, the two plausible species are either the Zwitter particle or the cation type of phenyl alanine. In conflict with the test results, the rate rule foretells a second-request rate dependence on $[H^+]$ with cation as the special species. The Zwitter particle is the dynamic species in this response, while protonated phenyl alanine isn't participated in the response succession.

It is realized that amino acids can be tracked down in the accompanying equilibria:




Figure 6: Perchloric acid's effects

It is possible that the protonation of PCC [Eq. (B)] is responsible for the damaging catalysis because it produces a more stable oxidant and an electrophile with open protonated and unprotonated structures. The development of protonated PCC types has remained undisclosed.



4. CONCLUSION

Within the sight of a low grouping of perchloric corrosive in a medium comprising of DMF and water, the response between amino acids and PCC is very languid when it happens at room temperature. When contrasted with [PCC], [Amino acid], and [HClO₄], the oxidation of amino corrosive by PCC is of the main request. The oxidation of the Zwitter ionic type of amino corrosive outcomes in the development of aldehyde. A few temperatures were explored to concentrate on the response. The legitimacy of the Arrhenius condition is laid out inside the temperature scope of 298-323 K. The way that the entropy esteem is negative focuses to the way that the perplexing C is more arranged than the reactant. The portrayal of the instrument concurs with each of the realities presented by the tests. The coordinated work on pyridinium chlorochromate oxidation of phenyl alanine in DMF-water media with perchloric destructive as a light source reveals that the unbiased amino destructive is involved in the reaction but that the

protonated amino destructive is unrelated to the cycle. It was finished at various temperatures generally through the reaction. The legitimacy of the Arrhenius condition is laid out inside the temperature scope of 298-323 K. The overall robotic succession that is introduced here is viable with the result of the item investigation as well as the discoveries of dynamic and unthinking exploration.

REFERENCES

1. Adari, K. K., Nowduri, A., & Vani, P. (2006). *Kinetics and mechanism of oxidation of l-Cysteine by Corey's reagent. Transition metal chemistry,*
2. Adari, K. K., Nowduri, A., & Vani, P. (2006). *Kinetics and mechanism of oxidation of l-Cysteine by Corey's reagent. Transition metal chemistry,*
3. Hawkins, C. L., & Davies, M. J. (2001). *Generation and propagation of radical reactions on proteins. Biochimica et Biophysica Acta (BBA)-Bioenergetics,*
4. Hiran, B. L., Meena, M. L., & Khuntwal, J. (2016). *Oxidation of phenyl alanine by pyridinium chlorochromate in acidic DMF–water medium: A kinetic study. Arabian Journal of Chemistry, 9, S1801-S1806.*
5. Jain, S., Hiran, B. L., & Bhatt, C. V. (2009). *Oxidation of some aliphatic alcohols by pyridinium chlorochromate-kinetics and mechanism. E-Journal of chemistry,*
6. Koohestani, B., Javanshir, Z., Ghammamy, S., Mehrani, K., Afrand, H., & Saghatforoush, L. (2008). *Synthesis and characterization of a new oxidation reagent: Tetrahexylammonium chlorochromate, (C₆H₁₃)₄N [CrO₃Cl]. Journal of the Mexican Chemical Society,*
7. Kumbhat, R., & Sharma, V. (2004). *Kinetics and mechanism of the oxidation of some unsaturated acids by pyridinium chlorochromate. Journal of the Indian Chemical Society,*
8. Mahanti, M.K., Banerji, K.K., (2004). *J. Indian Chem. Soc.*

9. Meena, M. L. (2023). *Kinetic and mechanistic study of oxidation of valine by pyridinium chlorochromate in aqueous DMF medium. World Journal of Biology Pharmacy and Health Sciences,*
10. Naik, P. N., Chimatadar, S. A., & Nandibewoor, S. T. (2008). *A kinetic and mechanistic study of the oxidation of tyrosine by chromium (VI) in aqueous perchloric acid medium. Transition Metal Chemistry,*
11. Nandibewoor, S.T., Naik, P.N., Chimatadar, S.A., (2008) *Transition Met. Chem.*
12. Patwari, S. B., Khansole, S. V., & Vibhute, Y. B. (2009). *Kinetics and mechanism of oxidation of aniline and substituted anilines by isoquinolinium bromochromate in aqueous acetic acid. Journal of the Iranian Chemical Society,*
13. Rajakumari, S., Chitradevi, S., Harikumar, S., Vivekanand, P. A., & Kamaraj, P. *Kinetics of oxidation and mechanism of alanine by pyridinium dichromate in the presence of perchloric acid medium in DMSO-H₂O (v/v) medium.*
14. Rathor, S., & Dangarh, B. K. *Kinetic and Mechanistic study of Oxidative Transformation of Some Amino acid by Pyridinium chlorochromate in DMF-Water Medium.*
15. REGAR, K. L., & Hiran, B. L. (2015). *Organic Catalysis in Oxidation of Isopropyl Alcohol by Pyridinium Fluorochromate-A Kinetic and Mechanistic Study. Chemical Science,*

Author's Declaration

I as an author of the above research paper/article, hereby, declare that the content of this paper is prepared by me and if any person having copyright issue or patent or anything otherwise related to the content, I shall always be legally responsible for any issue. For the reason of invisibility of my research paper on the website/amendments/updates, I have resubmitted my paper for publication on the same date. If any data or information given by me is not correct, I shall always be legally responsible. With my whole responsibility legally and formally have intimated the publisher (Publisher) that my paper has been checked by my guide (if any) or expert to make it sure that paper is technically right and there is no unaccepted plagiarism and hentriacontane is genuinely mine. If any issue arises related to Plagiarism/ Guide Name/ Educational Qualification/ Designation/ Address of my university/ college/institution/ Structure or Formatting/ Resubmission /Submission /Copyright /Patent/Submission for any higher degree or Job/Primary Data/Secondary Data Issues. I will be solely/entirely responsible for any legal issues. I have been informed that the most of the data from the website is invisible or shuffled or vanished from the database due to some technical fault or hacking and therefore the process of resubmission is there for the

scholars/students who finds trouble in getting their paper on the website. At the time of resubmission of my paper I take all the legal and formal responsibilities, If I hide or do not submit the copy of my original documents(Andhra/Driving License/Any Identity Proof and Photo) in spite of demand from the publisher then my paper maybe rejected or removed from the website anytime and may not be consider for verification. I accept the fact that As the content of this paper and the resubmission legal responsibilities and reasons are only mine then the Publisher (Airo International Journal/Airo National Research Journal) is never responsible. I also declare that if publisher finds Any complication or error or anything hidden or implemented otherwise, my paper maybe removed from the website or the watermark of remark/actuality maybe mentioned on my paper. Even if anything is found illegal publisher may also take legal action against me

Shobhag Singh Sisodiya
Dr. Rama Nand Yadav
